



ORIGINAL ARTICLE

WILEY

Hydrogen gas inhalation protects against cutaneous ischaemia/reperfusion injury in a mouse model of pressure ulcer

Wei Fang^{1,2,3}  | Guizhen Wang⁴ | Luyan Tang^{1,2} | Huilin Su¹ | Huyan Chen¹ | Wanqing Liao³ | Jinhua Xu^{1,2}

¹Department of Dermatology, Huashan Hospital, Fudan University, Shanghai, China

²The Shanghai Institute of Dermatology, Shanghai, China

³Department of Dermatology and Venereology, Changzheng Hospital, Shanghai Key Laboratory of Molecular Medical Mycology, Shanghai, China

⁴Emergency room, Shanghai Tenth People's Hospital of Tongji University, Shanghai, China

Correspondence

Jinhua Xu, Department of Dermatology, Huashan Hospital, Fudan University, Shanghai, China.
Email: xjhhsyy@163.com

Funding information

This study was supported by the Grant of Shanghai Key Medical Discipline for Dermatology (2017ZZ02002), the Grant of Shanghai leading talent project and China Postdoctoral Science Foundation Grant (2016M600286).

Abstract

Pressure ulcer formation depends on various factors among which repetitive ischaemia/reperfusion(I/R) injury plays a vital role. Molecular hydrogen (H₂) was reported to have protective effects on I/R injuries of various internal organs. In this study, we investigated the effects of H₂ inhalation on pressure ulcer and the underlying mechanisms. H₂ inhalation significantly reduced wound area, 8-oxo-dG level (oxidative DNA damage) and cell apoptosis rates in skin lesions. H₂ remarkably decreased ROS accumulation and enhanced antioxidant enzymes activities by up-regulating expression of Nrf2 and its downstream components in wound tissue and/or H₂O₂-treated endothelia. Meanwhile, H₂ inhibited the overexpression of MCP-1, E-selectin, P-selectin and ICAM-1 in oxidant-induced endothelia and reduced inflammatory cells infiltration and proinflammatory cytokines (TNF- α , IL-1, IL-6 and IL-8) production in the wound. Furthermore, H₂ promoted the expression of pro-healing factors (IL-22, TGF- β , VEGF and IGF1) and inhibited the production of MMP9 in wound tissue in parallel with acceleration of cutaneous collagen synthesis. Taken together, these data indicated that H₂ inhalation suppressed the formation of pressure ulcer in a mouse model. Molecular hydrogen has potentials as a novel and alternative therapy for severe pressure ulcer. The therapeutic effects of molecular hydrogen might be related to its antioxidant, anti-inflammatory, pro-healing actions.

KEYWORDS

hydrogen, inflammation, oxidative stress, pressure ulcer, reperfusion injury, wound healing

1 | INTRODUCTION

Pressure ulcer is a chronic inflammatory dermatosis primarily occurring over bony prominences (such as the sacrum, trochanter and the heels) of senior bedridden patients. Without prompt treatment, pressure ulcer may progress to cellulitis, osteomyelitis, sepsis and even

death. Pressure ulcer has become a serious public health problem owing to aging of the population.¹

There are several hypotheses for the mechanisms underlying chronic pressure ulcer, such as local tissue hypoxia, repetitive ischaemia/reperfusion (I/R), wound bacterial colonization, etc.^{2,3} Among these factors, I/R injury is thought to be a principal causative factor.^{4,5} Long-time recumbent position interrupts arteriolar capillary blood flow and thus induces local ischaemia in cutaneous tissue with

Wei Fang and Guizhen Wang contributed equally to this work.

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2018 The Authors. Journal of Cellular and Molecular Medicine published by John Wiley & Sons Ltd and Foundation for Cellular and Molecular Medicine.

prominent bony protrusions. After a change of body position, reperfusion of blood to the ischaemic skin initiates a series of harmful events because of a large increase in reactive oxygen species (ROS).⁶ Excessive ROS cause direct damage on lipids, proteins and nucleic acids, leading to cell apoptosis and tissue injury. Meanwhile, these free radicals induce the development of inflammatory responses such as endothelial dysfunction, neutrophil and macrophage infiltration, production of proinflammatory cytokines and, thereafter, tissues necrosis.^{6,7} Consistently, several studies demonstrated that skin ulcers induced by cutaneous I/R were inhibited by treatment with antioxidants such as vitamin E, melatonin and deferoxamine.⁸⁻¹⁰ Therefore, antioxidant administration may be a good therapeutic strategy for promoting healing of pressure ulcers.

In the past decade, H₂ as a novel medical gas has gained wide attention. In 2007, Ohsawa and colleagues found that H₂ affords neuroprotection against brain I/R injury by selectively neutralizing hydroxyl radicals and peroxynitrite.¹¹ Later studies demonstrated similar protective effects of hydrogen on I/R injuries of other organs such as liver, heart and intestines.¹²⁻¹⁴ The potential mechanisms might be involved in its antioxidant and anti-inflammatory and anti-apoptotic properties.¹⁵ These biological effects further enhanced the potential of H₂ in clinical application of various organ system diseases.^{15,16} Moreover, molecular hydrogen displayed high safety in vivo even at high concentration and high pressure, which could effectively reach target tissues and cells by gaseous diffusion but has no effect on physiological variables such as pH, oxygen saturation and blood pressure.^{17,18} Taken together, we hypothesized that H₂ could act as an effective treatment for pressure ulcer induced by repeated cutaneous I/R injury.

A recent study reported that hydrogen-water intake promoted wound size reduction and early recovery in 22 elderly in-patients with severe pressure ulcer.¹⁹ However, there has been no experimental evidence of the beneficial effects of molecular hydrogen on I/R-induced pressure ulcer using animal models. This study aimed to determine the possible protective effects of H₂ on pressure ulcer and the underlying mechanisms.

2 | MATERIALS AND METHODS

2.1 | Animals and cells

Female C57BL/6 mice (8 to 12 weeks old) were purchased from Changzhou Cavens Laboratory Animal Ltd. (Changzhou, China). All experiments were approved by the Ethical Committee for Animal Experiments of Fudan University, and strictly carried out in accordance with the approved guidelines. HUVEC cells were purchased from ATCC (Manassas, VA), and maintained in RPMI 1640 containing 10% foetal bovine serum (Thermo Fisher Scientific Inc., Waltham, MA).

2.2 | I/R cycles and analysis

The cutaneous I/R model was established according to previously published reports.²⁰⁻²² Briefly, all mice were anaesthetized, and their

backs were shaved and cleaned with 75% ethanol. The dorsal skin was gently pulled up and placed between two round ferrite magnetic plates that had a 12-mm diameter and 5-mm thickness (NeoMag Co, Ichikawa, Japan). A single I/R cycle was initiated with a 12-hour period of magnet placement, and followed by a release or rest period of 12 hours. After three I/R cycles, all of the mice developed two circular ulcers separated by a bridge of normal skin. For analysis, each wound site was digitally photographed after wounding, and wound areas were measured on photographs using Image J (version 1.48, NIH, Bethesda, MD) as previously described.^{20,23} To assess the effects of hydrogen gas on wound healing, the mice were housed in a specific airtight device producing air mixture including 2% or 75% H₂ and 21%O₂ for one week (6 hours per day) before the beginning of I/R cycles. H₂ treatment was continuously performed until the wounds completely healed. The hydrogen-producing device was provided by Shanghai Asclepius Meditec Co. Ltd (Shanghai, China). Wound sites were digitally photographed at various time-points after wounding, and wound areas were measured on the images using ImageJ software version 1.46r (NIH, Bethesda, MD).

2.3 | Histological and immunohistochemical examinations

The wounds were harvested with a 5-mm rim of unwounded skin tissue from sacrificed mice. Skin samples were fixed in 10% paraformaldehyde and embedded in paraffin. Sections (6 μm) were stained with HE and Masson or processed for subsequent immunostaining. For immunohistochemistry, deparaffinized sections were incubated with 3% H₂O₂ for 5 minutes to block endogenous peroxidase activity. After blocking with 10% foetal bovine serum, sections were stained with primary antibodies of interest followed by secondary Abs. Sections were washed three times with PBS buffer. The colour was developed using DAB substrate-chromogen solution (Bio-care Medical). The sections were then counterstained with HE.

2.4 | ROS measurement

ROS levels were measured in wound homogenates using ROS ELISA Kit (J&L Bio., Shanghai, China). For ROS detection in vitro, HUEVC cells were incubated in 2% or 75% H₂ incubator (Shanghai Asclepius Meditec Co. Ltd) for 24 hours and then stimulated with 0.25 mmol/L H₂O₂ (100 mL/well) for 2 hours as previous report.²³ The control group was without H₂ pretreatment. The ROS levels were then examined with dihydroethidium (DHE) (Beyotime, Shanghai, China) according to the manufacturer' protocol. Fluorescent pictures were collected by fluorescence microscope (Olympus IX71), and fluorescence intensity was analysed by Image J software (NIH, Bethesda, MD).

2.5 | CCK8 assay

The rate of cell proliferation was detected using CCK8 assay kit (Beyotime, Shanghai, China). After H₂ preincubation and 0.25 mmol/

L H₂O₂ treatment, 10⁴ of HUEVC cells were placed into 96-well plate (200 μ L per well) and 20 μ L CCK8 was added into each well for 24 hours. The optical density (OD) was read at 450 nm wavelength by ELISA microplate reader (Thermo Multiskan MK3).

2.6 | Apoptosis (TUNEL) assay

Apoptosis assay was carried out in both skin sections and H₂O₂-treated HUEVC cells using terminal deoxynucleotide transferase dUTP nick end labelling (TUNEL) staining kit (Roche Diagnostics, Indianapolis, IN) according to the manufacturer's instructions. Photographs were taken and visualized with inverted fluorescence microscope (Olympus IX71). The number of apoptotic cells was determined by counting TUNEL and Hoechst double positive nuclei in the field as previously described.²³

2.7 | Quantitative real-time PCR

For real-time PCR, total RNA was isolated from injured skin samples or H₂O₂-treated HUEVC cells using the Rneasy kit (QIAGEN Ltd., Crawley, UK) according to the manufacturer's protocols. First-strand cDNA was synthesized using the SuperScript III First-Strand Synthesis Kit (Invitrogen) according to its protocol. Resulting cDNA was used as template for subsequent real-time PCR using iQ SYBR Green Supermix (Bio-Rad) according to the manufacturer's recommendations. Primers (Table S1) were synthesized by Shanghai Sangon Biotech Co., Ltd (Shanghai, China). Relative expression of PCR products was determined using the 2^{- $\Delta\Delta$ CT} method and calculated relative to the control group.

2.8 | Western blot

For protein blotting, total proteins were extracted from homogenized skin tissue using as previously described.²⁴ Twenty micrograms of proteins was loaded into 8% SDS-PAGE and then transferred onto a polyvinylidene fluoride membrane. After blocking and washing, the membranes were incubated with the indicated primary Abs. The membranes were then incubated with horseradish peroxidase-labelled secondary antibody and developed with the ImmobilonTW Western Chemiluminescent HRP Substrate (Millipore, Billerica, MA). The blots were assessed by Image J software (NIH, Bethesda, MD).

All the antibodies (Abs) and their sources in the study are listed in Table S2.

2.9 | ELISA

Supernatants of mouse wound homogenates or cell culture medium were used for ELISA assay for SOD, GPx, CAT (BioVision), IL-1 β , TNF- α , IL-6, IL-8 and IL-22 (eBioscience) according to the manufacturers' recommendation. Total protein in the supernatant was detected with a commercial kit (BCA Protein Assay kit; Pierce, Rockford, IL). The data were expressed as anti-oxidative enzyme (unit/mL) or cytokine (pg/mL)/total protein (mg/mL) for each sample.

2.10 | Statistical analysis

Results were expressed as means \pm SEM. Unpaired, two-sided Student's *t* test was performed determine the statistical differences between the sample means using GraphPad Prism 6.0 (La Jolla, CA). *P* < .05 were considered statistically significant.

3 | RESULTS

3.1 | H₂ inhalation protected against cutaneous I/R-induced pressure ulcer formation

To assess the preventive effects of H₂ on the development of cutaneous pressure ulcers after I/R injury in vivo, a decubitus ulcer-like mouse model was constructed as previously described.^{20,23} In order to reach fully saturation in cutaneous tissue, 2% or 75% H₂ was inhaled by mice in experimental groups for one week (6 hours per day) before the beginning of I/R cycles according to the discovery of Scottish physiologist John Scott Haldane about the human body and the nature of gases.²⁵ Wound areas in H₂-treated mice were significantly smaller than those in control mice especially in early stage after I/R cycles (Figure 1). Inhalation of 75% H₂ significantly shortened the wound closure time in pressure ulcer mice. Furthermore, we also observed cleaner wounds and less scratching behaviour in H₂-treated mice than in control group. These results suggested that H₂ inhalation especially at high concentration protected the formation of cutaneous ulcers after I/R cycles.

3.2 | H₂ inhalation altered the histopathological characteristics of pressure ulcer skin after cutaneous I/R

Haematoxylin and eosin (HE) staining showed that H₂ inhalation reduced inflammatory cell infiltration and tissue necrosis in skin wound caused by I/R cycles (Figure S1). Compared with the control group, H₂-treatment groups (especially 75%H₂) displayed notable acceleration and enhancement in dermal collagen synthesis of skin wound (Figure 2). These results suggested that H₂ inhalation alleviated the inflammatory response and promoted the wound healing in pressure ulcer.

3.3 | H₂ inhalation alleviated oxidative DNA damage and suppressed apoptosis in skin tissues after cutaneous I/R

We further evaluated the levels of oxidative DNA damage (8-oxo-dG) and cell apoptosis after cutaneous I/R injury. Immunohistochemical (IHC) analysis showed that three groups had similar levels of 8-oxo-dG staining at 4 days postreperfusion. However, the area and intensity of positive staining in H₂-treated mice were significantly reduced compared to those in control mice and 75% H₂-treated group had lowest level of 8-oxo-dG at 6 and 8 days after reperfusion (Figure S2A). TUNEL analysis showed that wound tissues of




FIGURE 1 H₂ inhalation protected against ulcer formation in cutaneous I/R injury mice model. A, Representative photographs of wounds after cutaneous I/R in control or H₂-treated mice at 4, 6, 8, 10 and 13 days after reperfusion. B, Relative wound area after I/R injury in normal C57BL/6 mice with or without H₂ inhalation (N = 8 for each time point and groups). The ulcer size in control mice at 4 days after reperfusion was assigned a value of 100%. *P < .05; **P < .01; ***P < .001 compared to control




FIGURE 2 H₂ treatment promoted dermal collagen synthesis in skin wound caused by I/R cycles. Skin tissue samples from pressure ulcer mice of each group were fixed and paraffin embedded. The tissue sections were subjected to Masson staining. Representative images from the wound skin tissues are shown. Scale bar, 20 μ m

control mice had a large number of apoptotic cells, which was significantly reduced by H₂ inhalation (especially 75% H₂) at day 8 postreperfusion (Figure S2B and Figure 2C). These results suggested that H₂ inhalation could alleviate the oxidative DNA damage and suppress apoptosis of skin tissues, and high-concentration H₂ had optimal protective effect against cutaneous I/R injury.

3.4 | H₂ inhalation reduced ROS accumulation and up-regulated antioxidant enzyme activities in skin tissue after cutaneous I/R

As ROS is essential mediators of reperfusion induced tissue damage,²⁶⁻²⁹ we assessed ROS levels and the activities of major

antioxidant enzymes including superoxide dismutase (SOD), catalase (CAT) and glutathione peroxidase (GPx) in the wound tissue at day 6 postreperfusion. H₂ suppressed the ROS accumulation in skin tissue caused by I/R injury ($P < .01$) in a concentration-dependent manner (Figure 3A). Activities of SOD, CAT and GPx in mice inhaled H₂ were significantly higher than in control mice (Figure 3B-D).

3.5 | H₂ activated the NRF2-ARE pathway in skin tissue after cutaneous I/R

We next investigated whether NRF2/ARE antioxidant pathway mediated the inhibition of ROS level of pressure ulcer tissue by H₂. Quantitative real-time PCR analysis showed that H₂ significantly up-regulated the mRNA levels of NRF2 and its target genes (heme oxygenase-1, HO-1; NADH quinone oxidoreductase 1, NQO1; aldo-keto reductase, AKR1C1) in a concentration-dependent manner in the skin tissue (Figure 4A). Immunoblotting also confirmed the enhanced expression of NRF2-ARE pathway components in the skin tissue of H₂ treatment group (especially 75%) (Figure 4B). These results indicated that H₂ activated the NRF2 pathway in skin lesions after cutaneous I/R.

3.6 | H₂ inhalation inhibited cutaneous I/R-induced inflammation

As proinflammatory cytokines are important for tissue damage during I/R injury, the levels of proinflammatory cytokines in the wounded skin were assessed. At day 6 post-I/R, H₂ treatment significantly reduced the relative mRNA and protein expression of TNF- α ,

IL-1 β , IL-6 and IL-8 in wounded tissue, and their expression was lowest in pressure ulcer mice treated with 75% H₂ (Figure 5A and B). IHC further confirmed that the protein levels of TNF- α , IL-1 β , IL-6 and IL-8 in cutaneous I/R injured skin tissue of mice inhaled H₂ were substantially lower than those of control mice (Figure 5C). Meanwhile, H₂ treatment significantly up-regulated the expression of IL-22, an essential pro-healing cytokine involved in repair events on different models of epithelial regeneration,^{30,31} at both mRNA and protein levels and 75% H₂ treated group displayed the highest level of IL-22 (Figure 5), which was mainly expressed in the dermal layer of perilesional skin tissues in mice treated by H₂ (Figure 5C). These results indicated that hydrogen inhalation reversed proinflammatory effects of skin wound environment after cutaneous I/R.

3.7 | H₂ inhalation modulated the expression of healing-associated molecules in vivo after cutaneous I/R

The metalloprotease MMP9 and pro-healing factors TGF- β , VEGF and IGF-1 play crucial roles during proliferative and remodelling phases of wound healing. To investigate how H₂ improves wound healing in pressure ulcer mice, its effects on expression of aforementioned molecules were analysed by quantitative PCR and immunoblotting. H₂ inhalation significantly decreased the expression level of MMP9 and enhanced the expression of TGF- β 1, VEGF and IGF-1 at both mRNA and protein levels in the wound. Mice treated with 75% H₂ had the lowest level of MMP9 and highest expression of pro-healing factors in skin tissue among three groups (Figure 6).




FIGURE 3 H₂ inhalation reduced ROS accumulation and enhanced antioxidant enzyme activities in skin tissue. ELISA assays for detection of ROS production (A) and the activities of SOD (B), GPx (C), and CAT (D). * $P < .05$; ** $P < .01$ compared to control




FIGURE 4 Hydrogen activated the NRF2-ARE pathway in wounded tissue. A, The mRNA levels of NRF2, HO-1, NQO1 and AKR1C1 were analysed by quantitative RT-PCR assay. B, Immunoblots of NRF2, HO-1, NQO1, AKR1C1 and β -actin. * $P < .05$; ** $P < .01$ compared to control

3.8 | H₂ reversed the adverse effect of H₂O₂ on cell proliferation and apoptosis of vascular endothelial cells

I/R injury is closely associated with microvascular dysfunction, which is largely a consequence of endothelial cell impairment.⁷ Thus, we further examine the effects of H₂ on human umbilical vein endothelial cells (HUVECs) against oxidative damage. Preincubation with H₂ for 24 hours significantly relieved the 0.25 mmol/L H₂O₂-induced proliferation inhibition of HUVECs in a concentration-dependent manner ($P < .01$) (Figure S3A). After exposure to 0.25 mmol/L H₂O₂, HUVECs showed a 2.5-folds increase in the apoptosis rate, which was markedly reduced by H₂ pretreatment (especially 75% H₂) ($P < .01$) (Figure S3B and C).

3.9 | H₂ rectified the imbalance of oxidation-reduction system in H₂O₂-treated vascular endothelial cells through the activation of Nrf2/ARE pathway

To further understand the mechanism of H₂-mediated protective effects on endothelial cells during cutaneous I/R, a series of in vitro studies were performed in oxidant-treated HUVECs. H₂O₂ notably up-regulated intracellular ROS production (Figure S4A) and suppressed the expression of various anti-oxidative enzymes (SOD, GPx, and CAT) (Figure S4B-D) in HUVECs, suggesting of a severe disturbance of intracellular oxidation-reduction system. Preincubation with H₂ resulted in a trend towards normalization of both intracellular ROS and anti-oxidative enzyme in H₂O₂-treated HUVECs (Figure S4A-D). Quantitative PCR assay demonstrated that H₂ significantly up-regulated the expression of NRF2 and its target genes

HO-1, AKR1C1 and NQO1 in HUVECs exposed to H₂O₂ in a concentration-dependent manner (Figure S4E-H). Together, these results suggested that H₂ might maintain intracellular homeostasis of oxidation-reduction system in vascular endothelial cells through the activation of Nrf2/ARE pathway.

3.10 | H₂ inhibited the expression of several chemokine and adhesion molecules in vascular endothelial cells stimulated by H₂O₂

Endothelial damage is a critical event in the early phase of inflammatory responses induced by cutaneous I/R, which could drive the up-regulation of various chemokines and adhesion molecules and then mediate leucocyte recruit, adhesion and emigration.^{7,32} Vascular endothelial cells more than doubled the transcriptional levels of MCP-1 (monocyte chemoattractant protein-1), E-selectin, P-selectin and ICAM-1 (intercellular cell adhesion molecule-1) after H₂O₂ exposure (Figure 7). Pretreatment with H₂ significantly suppressed the elevation of their expression in H₂O₂-exposed HUVECs ($P < .01$) (Figure 7). The inhibitory effects were positively correlated with H₂ concentration. Our results suggested that H₂ might exert anti-inflammatory effects against cutaneous I/R injury by down-regulating the expression of endothelial chemokines and adhesion molecules.

4 | DISCUSSION

In the present study, we first demonstrated the protective effects of H₂ inhalation on pressure ulcers in a murine model. H₂ treatment significantly reduced the wound area in a concentration-dependent




FIGURE 5 Hydrogen altered the expression or production of inflammatory cytokines during IR cycles. A, mRNA expression of TNF- α , IL-1 β , IL-6, IL-8 and IL-22 in skin wounds. B, Immunoblot detecting the protein levels of TNF- α , IL-1 β , IL-6, IL-8 and IL-22 in wounded skin tissue. C, IHC assessing injured skin TNF- α , IL-1 β , IL-6, IL-8 and IL-22 levels. * $P < .05$; ** $P < .01$ compared to control

manner in the early phase after cutaneous I/R. The improvement in skin ulcer recovery induced by H₂ was paralleled by significant reductions in oxidative DNA damage, cell apoptosis and acceleration of cutaneous collagen synthesis. This result was consistent with a clinical study that hydrogen-water intake (0.8-1.3 ppm, 600 mL per day) via tube-feeding significantly reduced the wound size in 22 hospitalized patients with pressure ulcer.¹⁹ Theoretically, hydrogen administration through respiratory route displays higher blood concentration and more accessibility for cutaneous lesions than those through gastrointestinal absorption.¹⁵ Therefore, H₂ inhalation has a greater potential to be applied in the prevention and treatment of patients with pressure ulcer.

H₂-mediated protection against pressure ulcers was primarily dependent on its antioxidant property. Oxidative stress plays

crucial roles in the initiation and progression phases of cutaneous I/R injury.^{3,6,26} In pressure ulcers, blood reperfusion after cutaneous ischaemia induced abundant reactive oxygen species (ROS), which could directly damage skin tissue and cause cell apoptosis. These toxic free radicals further aggravated leucocyte infiltration, which in turn triggered more ROS release and thus trapped in a vicious circle. In our study, H₂ treatment remarkably decreased the ROS accumulation and 8-oxo-dG formation (a sensitive biomarker of oxidative DNA damage), and also enhanced the activities of several antioxidant enzymes (such as SOD, GPx and CAT) in wound tissue and/or H₂O₂-treated HUEVCs. The antioxidant ability was positively correlated with its dosage, and high concentration of H₂ was more advantageous in maintaining intracellular




FIGURE 6 Hydrogen modulated the expression of healing-associated molecules in skin wounds. A, mRNA expression of MMP9, TGF- β , VEGF and IGF-1 was analysed by quantitative RT-PCR. B, Immunoblot assessment of MMP9, TGF- β , VEGF and IGF-1 protein levels in wounded skin tissue. * $P < .05$; ** $P < .01$ compared to control




FIGURE 7 Hydrogen decreased the expression of several molecules associated with leucocyte-endothelium in H₂O₂-treated HUEVCs. Quantitative RT-PCR assay of MCP-1 (A), P-selectin (B), E-selectin (C) and ICAM-1(D) in HUEVCs. ** $P < .01$, * $P < .05$, compared to the control group; ## $P < .01$, # $P < .05$, compared to H₂O₂-treated group

homeostasis of oxidation-reduction system in skin tissue after I/R injury.

We also determined the mechanism underlying the antioxidant property of H₂ against cutaneous I/R injury, which was closely associated with the activation of Nrf2/ARE pathway. The transcription factor Nrf2 is essential for regulating the adaptive response to exogenous and endogenous oxidative stresses.^{33,34} Under moderate oxidative stress, Nrf2 translocates to the nucleus where it binds to ARE and induces the transcription of downstream antioxidant genes. In our study, H₂ treatment significantly up-regulated the expression of Nrf2 and its downstream targets such as HO-1, NQO1 and AKR1C1 in wound tissue and HUEVCs exposed to H₂O₂. Similar phenomena

were also observed in previous studies of other organ I/R injury or other inflammatory diseases.³⁵⁻³⁸ Molecular hydrogen could attenuate intestinal injury in wild-type but not Nrf2-knockout mice with severe sepsis by regulating HO-1 expression.³⁷ Therefore, hydrogen might activate the Nrf2/ARE pathway to restore the homeostasis of cutaneous oxidation-reduction system against I/R injury.

The protective effects of H₂ were also attributed to its potent anti-inflammatory activity. After I/R cycles, plenty of macrophages and neutrophils were accumulated in the treated tissue accompanied by significant production of proinflammatory cytokines, which caused skin injury and necrosis in the wound.²¹ In our study, H₂ inhalation attenuated inflammatory cells infiltration and cutaneous

necrosis in the wound induced by I/R injury. Proinflammatory cytokines such as TNF- α , IL-1, IL-6 and IL-8 were decreased while IL-22 was increased in the wounded skin of H₂-treated mice. Distinct from proinflammatory cytokines, IL-22 mediates a crosstalk between immune system and cutaneous cells (such as fibroblasts and keratinocytes) and plays a pro-healing role in wound repair.^{30,39} Our data demonstrated that H₂ reversed the excessive inflammatory response induced by cutaneous I/R injury.

The anti-inflammatory property of hydrogen might be partially attributed to its protection on vascular endothelial cells against oxidative stress. During I/R injury, endothelial damage initiated leucocyte adhesion, recruitment and infiltration by up-regulating the expression of chemokines and adhesion molecules.^{6,7,32} Our in vitro assay showed that H₂ treatment significantly alleviated the oxidative injury of endothelial cells and reduced the overexpression of MCP-1, E-selectin, P-selectin and ICAM-1 in HUEVCs exposed to H₂O₂. MCP-1 is a critical molecule for chemotaxis and activation of macrophage, which is a significant source of proinflammatory cytokines and contributes to I/R injury of skin and other organs.^{21,40-42} The adhesion molecules studied here were mainly responsible for leucocyte rolling, localization and adhesion to the endothelium.^{43,44} Hence, hydrogen might modulate those molecules associated with leucocyte-endothelium interaction to inhibit subsequent inflammatory reaction caused by cutaneous I/R injury.

In pressure ulcer, the wound sites were constantly in the dynamic pathological alterations of inflammatory injury and tissue repair. Outcome of wound healing process was mainly determined by the presence and concentration of the healing-associated factors such as MMPs, TGF- β , VEGF and IGF1.⁴⁵⁻⁴⁷ Overproduction or high activity of MMP9 and suppressed expression of TGF- β were identified as indicators of poor healing in skin samples of chronic ulcers.^{46,48} In our study, H₂ promoted the expression of pro-healing factors (TGF- β , VEGF and IGF1) and inhibited the production of MMP9 in wound tissue of pressure ulcer, accompanied by acceleration of dermal collagen synthesis. Hence, hydrogen had a wound healing promoting effect against cutaneous I/R injury.

Taken together, the present results indicated that hydrogen suppressed the formation of decubitus ulcers by its antioxidant, anti-inflammatory, pro-healing activities against cutaneous I/R injury. Similar therapeutic effects of hydrogen were also reported in common senile diseases concomitant with decubitus, such as cerebral or myocardial infarctions, COPD, diabetes, hyperlipaemia, malignant tumours.^{11,36,49-54} Furthermore, H₂ had no cytotoxicity *in vivo* in human body even at a high concentration.¹⁷ Therefore, hydrogen gas has a great potential for preventing and/or treating pressure ulcer.

There were still some limitations in our study. Firstly, we did not detect the cutaneous hydrogen concentration, which might provide direct evidence to support the dosage-dependent protection of hydrogen gas against cutaneous I/R injury. In addition, the exact mechanisms by which hydrogen modulates oxidative stress, inflammation and wound repair in pressure ulcer were still unclear. More experiments are needed to work out these problems before the clinical trial of H₂ in pressure ulcers.

ACKNOWLEDGEMENTS

We extend our sincerest gratitude to the staff of Shanghai Asclepius Meditec Co., Ltd. for their vital technical support in the hydrogen-oxygen nebulizer device. We also thank Prof. Zhimin Kang of Shanghai Huikang Hydrogen Medical Research Center for technical assistance in study design.

CONFLICT OF INTEREST

The authors have no conflicts of interest to report.

ORCID

Wei Fang  <http://orcid.org/0000-0002-0749-298X>

REFERENCES

- Gorecki C, Brown JM, Nelson EA, et al. Impact of pressure ulcers on quality of life in older patients: a systematic review. *J Am Geriatr Soc.* 2009;57:1175-1183.
- Mustoe T. Understanding chronic wounds: a unifying hypothesis on their pathogenesis and implications for therapy. *Am J Surg.* 2004;187:655-705.
- Mustoe TA, O'Shaughnessy K, Kloeters O. Chronic wound pathogenesis and current treatment strategies: a unifying hypothesis. *Plast Reconstr Surg.* 2006;117:35S-41S.
- Salcido R, Donofrio JC, Fisher SB, et al. Histopathology of pressure ulcers as a result of sequential computer-controlled pressure sessions in a fuzzy rat model. *Adv Wound Care.* 1994;7:23-24, 26, 28 passim.
- Peirce SM, Skalak TC, Rodeheaver GT. Ischemia-reperfusion injury in chronic pressure ulcer formation: a skin model in the rat. *Wound Repair Regen.* 2000;8:68-76.
- Eltzschig HK, Collard CD. Vascular ischaemia and reperfusion injury. *Br Med Bull.* 2004;70:71-86.
- Seal JB, Gewertz BL. Vascular dysfunction in ischemia-reperfusion injury. *Ann Vasc Surg.* 2005;19:572-584.
- Houwing R, Overgoor M, Kon M, Jansen G, van Asbeck BS, Haalboom JR. Pressure-induced skin lesions in pigs: reperfusion injury and the effects of vitamin E. *J Wound Care.* 2000;9:36-40.
- Sener G, Sert G, Ozer SA, Arbak S, Gedik N, Ayanoglu-Duylger G. Melatonin protects against pressure ulcer-induced oxidative injury of the skin and remote organs in rats. *J Pineal Res.* 2006;40:280-287.
- Duscher D, Neofytou E, Wong VW, et al. Transdermal deferoxamine prevents pressure-induced diabetic ulcers. *Proc Natl Acad Sci U S A.* 2015;112:94-99.
- Ohsawa I, Ishikawa M, Takahashi K, et al. Hydrogen acts as a therapeutic antioxidant by selectively reducing cytotoxic oxygen radicals. *Nat Med.* 2007;13:688-694.
- Liu Q, Shen WF, Sun HY, et al. Hydrogen-rich saline protects against liver injury in rats with obstructive jaundice. *Liver Int.* 2010;30:958-968.
- Sun Q, Kang Z, Cai J, et al. Hydrogen-rich saline protects myocardium against ischemia/reperfusion injury in rats. *Exp Biol Med (Maywood).* 2009;234:1212-1219.
- Zheng X, Mao Y, Cai J, et al. Hydrogen-rich saline protects against intestinal ischemia/reperfusion injury in rats. *Free Radic Res.* 2009;43:478-484.
- Ge L, Yang M, Yang NN, Yin XX, Song WG. Molecular hydrogen: a preventive and therapeutic medical gas for various diseases. *Oncotarget.* 2017;8:102653-102673.

16. Ostojic SM. Molecular hydrogen: an inert gas turns clinically effective. *Ann Med*. 2015;47:301-304.
17. Abraini JH, Gardette-Chauffour MC, Martinez E, Rostain JC, Lemaire C. Psychophysiological reactions in humans during an open sea dive to 500 m with a hydrogen-helium-oxygen mixture. *J Appl Physiol*. 1994;76:1113-1118.
18. Ono H, Nishijima Y, Adachi N, et al. A basic study on molecular hydrogen (H₂) inhalation in acute cerebral ischemia patients for safety check with physiological parameters and measurement of blood H₂ level. *Med Gas Res*. 2012;2:21.
19. Li Q, Kato S, Matsuoka D, Tanaka H, Miwa N. Hydrogen water intake via tube-feeding for patients with pressure ulcer and its reconstructive effects on normal human skin cells in vitro. *Med Gas Res*. 2013;3:20.
20. Uchiyama A, Yamada K, Perera B, et al. Protective effect of MFG-E8 after cutaneous ischemia-reperfusion injury. *J Invest Dermatol*. 2015;135:1157-1165.
21. Saito Y, Hasegawa M, Fujimoto M, et al. The loss of MCP-1 attenuates cutaneous ischemia-reperfusion injury in a mouse model of pressure ulcer. *J Invest Dermatol*. 2008;128:1838-1851.
22. Stadler I, Zhang RY, Oskoui P, Whittaker MS, Lanzafame RJ. Development of a simple, noninvasive, clinically relevant model of pressure ulcers in the mouse. *J Invest Surg*. 2004;17:221-227.
23. Uchiyama A, Yamada K, Perera B, et al. Protective effect of botulinum toxin A after cutaneous ischemia-reperfusion injury. *Sci Rep*. 2015;5:9072.
24. Long M, de la Vega MR, Wen Q, et al. An essential role of NRF2 in diabetic wound healing. *Diabetes*. 2016;65:780-793.
25. Bove A, Davis J. *Bove and davis' diving medicine* (4th edn.). New York, NY: Saunders; 2003:53-76.
26. Siemionow M, Arslan E. Ischemia/reperfusion injury: a review in relation to free tissue transfers. *Microsurgery*. 2004;24:468-475.
27. Chouchani ET, Pell VR, James AM, et al. A unifying mechanism for mitochondrial superoxide production during ischemia-reperfusion injury. *Cell Metab*. 2016;23:254-263.
28. Minutoli L, Puzzolo D, Rinaldi M, et al. ROS-mediated NLRP3 inflammasome activation in brain, heart, kidney, and testis ischemia/reperfusion injury. *Oxid Med Cell Longev*. 2016;2016:2183026.
29. Granger DN, Kvietys PR. Reperfusion injury and reactive oxygen species: the evolution of a concept. *Redox Biol*. 2015;6:524-551.
30. Avitabile S, Odoriso T, Madonna S, et al. Interleukin-22 promotes wound repair in diabetes by improving keratinocyte pro-healing functions. *J Invest Dermatol*. 2015;135:2862-2870.
31. Pickert G, Neufert C, Leppkes M, et al. STAT3 links IL-22 signaling in intestinal epithelial cells to mucosal wound healing. *J Exp Med*. 2009;206:1465-1472.
32. Toledo-Pereyra LH, Toledo AH, Walsh J, Lopez-Nebolina F. Molecular signaling pathways in ischemia/reperfusion. *Exp Clin Transplant*. 2004;2:174-177.
33. Kensler TW, Wakabayashi N, Biswal S. Cell survival responses to environmental stresses via the Keap1-Nrf2-ARE pathway. *Annu Rev Pharmacol Toxicol*. 2007;47:89-116.
34. Jaramillo MC, Zhang DD. The emerging role of the Nrf2-Keap1 signaling pathway in cancer. *Genes Dev*. 2013;27:2179-2191.
35. Yu J, Zhang W, Zhang R, et al. Molecular hydrogen attenuates hypoxia/reoxygenation injury of intrahepatic cholangiocytes by activating Nrf2 expression. *Toxicol Lett*. 2015;238:11-19.
36. Song G, Zong C, Zhang Z, et al. Molecular hydrogen stabilizes atherosclerotic plaque in low-density lipoprotein receptor-knockout mice. *Free Radic Biol Med*. 2015;87:58-68.
37. Yu Y, Yang Y, Bian Y, et al. Hydrogen gas protects against intestinal injury in wild type but not NRF2 knockout mice with severe sepsis by regulating HO-1 and HMGB1 release. *Shock*. 2017;48:364-370.
38. Tamaki N, Orihuela-Campos RC, Fukui M, Ito HO. Hydrogen-rich water intake accelerates oral palatal wound healing via activation of the Nrf2/antioxidant defense pathways in a rat model. *Oxid Med Cell Longev*. 2016;2016:5679040.
39. McGee HM, Schmidt BA, Booth CJ, et al. IL-22 promotes fibroblast-mediated wound repair in the skin. *J Invest Dermatol*. 2012;133:1321-1329.
40. Leonard EJ, Yoshimura T. Human monocyte chemoattractant protein-1 (MCP-1). *Immunol Today*. 1990;11:97-101.
41. Hayasaki T, Kaikita K, Okuma T, et al. CC chemokine receptor-2 deficiency attenuates oxidative stress and infarct size caused by myocardial ischemia-reperfusion in mice. *Circ J*. 2006;70:342-351.
42. Furuichi K, Wada T, Iwata Y, et al. CCR2 signaling contributes to ischemia-reperfusion injury in kidney. *J Am Soc Nephrol*. 2003;14:2503-2515.
43. Sawaya DE, Zibari GB, Minardi A, et al. P-selectin contributes to the initial recruitment of rolling and adherent leukocytes in hepatic venules after ischemia/reperfusion. *Shock*. 1999;12:227-232.
44. Kurose I, Anderson DC, Miyasaka M, et al. Molecular determinants of reperfusion-induced leukocyte adhesion and vascular protein leakage. *Circ Res*. 1994;74:336-343.
45. Ravanti L, Kähäri VM. Matrix metalloproteinases in wound repair (review). *Int J Mol Med*. 2000;6:391-407.
46. Pakyari M, Farrokhi A, Maharlooie MK, Ghahary A. Critical role of transforming growth factor beta in different phases of wound healing. *Adv Wound Care (New Rochelle)*. 2013;2:215-224.
47. Barrientos S, Stojadinovic O, Golinko MS, Brem H, Tomic-Canic M. Growth factors and cytokines in wound healing. *Wound Repair Regen*. 2008;16:585-601.
48. Yager DR, Zhang LY, Liang HX, Diegelmann RF, Cohen IK. Wound fluids from human pressure ulcers contain elevated matrix metalloproteinase levels and activity compared to surgical wound fluids. *J Invest Dermatol*. 1996;107:743-748.
49. Liu SL, Liu K, Sun Q, Liu WW, Tao HY, Sun XJ. Hydrogen therapy may be a novel and effective treatment for COPD. *Front Pharmacol*. 2011;2:19.
50. Kawai D, Takaki A, Nakatsuka A, et al. Hydrogen-rich water prevents progression of nonalcoholic steatohepatitis and accompanying hepatocarcinogenesis in mice. *Hepatology*. 2012;56:912-921.
51. Dole M, Wilson FR, Fife WP. Hyperbaric hydrogen therapy: a possible treatment for cancer. *Science*. 1975;190:152-154.
52. Zong C, Song G, Yao S, et al. Administration of hydrogen-saturated saline decreases plasma low-density lipoprotein cholesterol levels and improves high-density lipoprotein function in high-fat diet-fed hamsters. *Metabolism*. 2012;61:794-800.
53. Hayashida K, Sano M, Ohsawa I, et al. Inhalation of hydrogen gas reduces infarct size in the rat model of myocardial ischemia-reperfusion injury. *Biochem Biophys Res Commun*. 2008;373:30-35.
54. Kamimura N, Nishimaki K, Ohsawa I, Ohta S. Molecular hydrogen improves obesity and diabetes by inducing hepatic FGF21 and stimulating energy metabolism in db/db mice. *Obesity (Silver Spring)*. 2011;19:1396-1403.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of the article.

How to cite this article: Fang W, Wang G, Tang L, et al. Hydrogen gas inhalation protects against cutaneous ischaemia/reperfusion injury in a mouse model of pressure ulcer. *J Cell Mol Med*. 2018;00:1-10. <https://doi.org/10.1111/jcmm.13704>